

Exam 3 : March 31

HW 10, Part I

Question 4 of 8

Compute T_3 for $f(x) = 3\sqrt{x}$ centered at $a = 1$.

(Use symbolic notation and fractions where needed.)

$$T_3(x) = \frac{3}{0!}(x-1)^0 + \frac{\frac{3}{2}}{1!}(x-1)^1 + \frac{-\frac{3}{4}}{2!}(x-1)^2 + \frac{\frac{9}{8}}{3!}(x-1)^3$$

$$= 3 + \frac{3}{2}(x-1) - \frac{3}{8}(x-1)^2 + \frac{9}{8 \cdot 6}(x-1)^3$$

$$= 3 + \frac{3}{2}(x-1) - \frac{3}{8}(x-1)^2 + \frac{3}{16}(x-1)^3$$

$$T(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} \cdot (x-a)^n$$

$$f(x) = 3\sqrt{x} \quad f(1) = 3$$

$$f'(x) = \frac{3}{2}x^{-1/2} \quad f'(1) = \frac{3}{2}$$

$$f''(x) = -\frac{3}{4}x^{-3/2} \quad f''(1) = -\frac{3}{4}$$

$$f'''(x) = \frac{9}{8}x^{-5/2} \quad f'''(1) = \frac{9}{8}$$

function

Use a plot of the error $|f(x) - T_3(x)|$ to find the largest value $c > 1$ such that the error on the interval $[1, c]$ is at most 0.75.

(Use decimal notation. Give your answer to three decimal places.)

c =

$$E(x) = \left| 3\sqrt{x} - \left[3 + \frac{3}{2}(x-1) - \frac{3}{8}(x-1)^2 + \frac{3}{16}(x-1)^3 \right] \right|$$

HW Part 4 #6

Express the definite integral as an infinite series in the form $\sum_{n=0}^{\infty} a_n$.

$$\int_0^1 2 \tan^{-1}(x^2) dx$$

(Express numbers in exact form. Use symbolic notation and fractions where needed.)

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + x^4 + \dots$$

↓ sub ($x \rightarrow -x^2$)

$$\frac{1}{1+x^2} = 1 - x^2 + x^4 - x^6 + x^8 - \dots$$

↓ integrate

$$\tan^{-1}x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \frac{x^9}{9} - \dots$$

$a_n =$

$$\tan^{-1}x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \frac{x^9}{9} - \dots = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1} \quad \text{@ } x \rightarrow x^2 \quad \sum_{n=0}^{\infty} (-1)^n \frac{(x^2)^{2n+1}}{2n+1}$$

$$\tan^{-1}(x^2) = x^2 - \frac{x^6}{3} + \frac{x^{10}}{5} - \frac{x^{14}}{7} + \frac{x^{18}}{9} - \dots = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+2}}{2n+1}$$


$\begin{matrix} | & | & | \\ n=0 & n=1 & n=2 \end{matrix}$

$k=4 \left\{ \begin{array}{l} kn+2=2 \leftrightarrow 0=n \\ kn+2=6 \leftrightarrow 1=n \\ kn+2=10 \leftrightarrow 2=n \end{array} \right.$

$\left(\int_0^1 x^2 \right) - \left(\int_0^1 \frac{x^6}{3} \right) = 2 \left[\frac{x^3}{3} - \frac{x^7}{7} \right] \Big|_0^1$

$$\begin{aligned} \int_0^1 2 \tan^{-1}x dx &= \int_0^1 2 \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+2}}{2n+1} dx = 2 \int_0^1 \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+2}}{2n+1} dx = 2 \sum_{n=0}^{\infty} \int_0^1 (-1)^n \frac{x^{4n+2}}{2n+1} dx \\ &= 2 \sum_{n=0}^{\infty} (-1)^n \int_0^1 \frac{x^{4n+2}}{2n+1} dx = 2 \sum_{n=0}^{\infty} (-1)^n \frac{1}{2n+1} \int_0^1 \frac{x^{4n+2}}{x^{4n+3}} dx = 2 \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+3}}{(2n+1)(4n+3)} \Big|_0^1 \\ &= 2 \sum_{n=0}^{\infty} (-1)^n \frac{1}{(2n+1)(4n+3)} - \frac{0}{(2n+1)(4n+3)} = \sum_{n=0}^{\infty} (-1)^n \frac{2}{(2n+1)(4n+3)} \end{aligned}$$

Error incurred by approximating using only n terms of an alternating series is less than the (n+1) term



this is an alt. series, which are easy to approximate!

$$\text{sub } (n+1) \quad \frac{2}{(2(n+1)+1)(4(n+1)+3)} = \frac{2}{(2n+3)(4n+7)} < 10^{-5}$$

set/solve this $\frac{2}{(2n+3)(4n+7)} < 10^{-5}$

$$200000 = 2 \cdot 10^5 = \frac{2}{10^{-5}} < (2n+3)(4n+7) = 8n^2 + 26n + 21$$

say @ $n=100$ this happens!

$$\int_0^1 2 \tan^{-1}x = \sum_{n=0}^{100} (-1)^n \frac{2}{(2n+1)(4n+3)}$$